

DIPHTHERIA TETANUS & PERTUSSIS VACCINES

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 Why get vaccinated?

Diphtheria, tetanus, and pertussis are serious diseases caused by bacteria. Diphtheria and pertussis are spread from person to person. Tetanus enters the body through cuts or wounds.

DIPHTHERIA causes a thick covering in the back of the throat.

- It can lead to breathing problems, paralysis, heart failure, and even death.

TETANUS (Lockjaw) causes painful tightening of the muscles, usually all over the body.

- It can lead to “locking” of the jaw so the victim cannot open his mouth or swallow. Tetanus leads to death in up to 2 out of 10 cases.

PERTUSSIS (Whooping Cough) causes coughing spells so bad that it is hard for infants to eat, drink, or breathe. These spells can last for weeks.

- It can lead to pneumonia, seizures (jerking and staring spells), brain damage, and death.

Diphtheria, tetanus, and pertussis vaccine (DTaP) can help prevent these diseases. Most children who are vaccinated with DTaP will be protected throughout childhood. Many more children would get these diseases if we stopped vaccinating.

DTaP is a safer version of an older vaccine called DTP. DTP is no longer used in the United States.

2 Who should get DTaP vaccine and when?

Children should get 5 doses of DTaP vaccine, one dose at each of the following ages:

- ✓ 2 months
- ✓ 4 months
- ✓ 6 months
- ✓ 15-18 months
- ✓ 4-6 years

DTaP may be given at the same time as other vaccines.

3

Some children should not get DTaP vaccine or should wait

- Children with minor illnesses, such as a cold, may be vaccinated. But children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.
- Any child who had a life-threatening allergic reaction after a dose of DTaP should not get another dose.
- Any child who suffered a brain or nervous system disease within 7 days after a dose of DTaP should not get another dose.
- Talk with your doctor if your child:
 - had a seizure or collapsed after a dose of DTaP,
 - cried non-stop for 3 hours or more after a dose of DTaP,
 - had a fever over 105°F after a dose of DTaP.

Ask your health care provider for more information. Some of these children should not get another dose of pertussis vaccine, but may get a vaccine without pertussis, called **DT**.

4

Older children and adults

DTaP is not licensed for adolescents, adults, or children 7 years of age and older.

But older people still need protection. A vaccine called **Tdap** is similar to DTaP. A single dose of Tdap is recommended for people 11 through 64 years of age. Another vaccine, called **Td**, protects against tetanus and diphtheria, but not pertussis. It is recommended every 10 years. There are separate Vaccine Information Statements for these vaccines.

5 What are the risks from DTaP vaccine?

Getting diphtheria, tetanus, or pertussis disease is much riskier than getting DTaP vaccine.

However, a vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of DTaP vaccine causing serious harm, or death, is extremely small.

Mild Problems (Common)

- Fever (up to about 1 child in 4)
- Redness or swelling where the shot was given (up to about 1 child in 4)
- Soreness or tenderness where the shot was given (up to about 1 child in 4)

These problems occur more often after the 4th and 5th doses of the DTaP series than after earlier doses. Sometimes the 4th or 5th dose of DTaP vaccine is followed by swelling of the entire arm or leg in which the shot was given, lasting 1-7 days (up to about 1 child in 30).

Other mild problems include:

- Fussiness (up to about 1 child in 3)
- Tiredness or poor appetite (up to about 1 child in 10)
- Vomiting (up to about 1 child in 50)

These problems generally occur 1-3 days after the shot.

Moderate Problems (Uncommon)

- Seizure (jerking or staring) (about 1 child out of 14,000)
- Non-stop crying, for 3 hours or more (up to about 1 child out of 1,000)
- High fever, over 105°F (about 1 child out of 16,000)

Severe Problems (Very Rare)

- Serious allergic reaction (less than 1 out of a million doses)
- Several other severe problems have been reported after DTaP vaccine. These include:
 - Long-term seizures, coma, or lowered consciousness
 - Permanent brain damage.

These are so rare it is hard to tell if they are caused by the vaccine.

Controlling fever is especially important for children who have had seizures, for any reason. It is also important if another family member has had seizures. You can reduce fever and pain by giving your child an *aspirin-free* pain reliever when the shot is given, and for the next 24 hours, following the package instructions.

6 What if there is a moderate or severe reaction?

What should I look for?

Any unusual conditions, such as a serious allergic reaction, high fever or unusual behavior. Serious allergic reactions are extremely rare with any vaccine. If one were to occur, it would most likely be within a few minutes to a few hours after the shot. Signs can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness. If a high fever or seizure were to occur, it would usually be within a week after the shot.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not provide medical advice

7 The National Vaccine Injury Compensation Program

In the rare event that you or your child has a serious reaction to a vaccine, a federal program has been created to help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit the program's website at www.hrsa.gov/vaccinecompensation.

8 How can I learn more?

- Ask your health care provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department's immunization program.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)**
 - Visit the National Immunization Program's website at www.cdc.gov/vaccines



U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Disease Control and Prevention

Polio Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages.
See www.immunize.org/vis.

Hojas de Información Sobre Vacunas están disponibles en español y en muchos otros idiomas. Visite <http://www.immunize.org/vis>

1 What is polio?

Polio is a disease caused by a virus. It enters the body through the mouth. Usually it does not cause serious illness. But sometimes it causes paralysis (can't move arm or leg), and it can cause meningitis (irritation of the lining of the brain). It can kill people who get it, usually by paralyzing the muscles that help them breathe.

Polio used to be very common in the United States. It paralyzed and killed thousands of people a year before we had a vaccine.

2 Why get vaccinated?

Inactivated Polio Vaccine (IPV) can prevent polio.

History: A 1916 polio epidemic in the United States killed 6,000 people and paralyzed 27,000 more. In the early 1950's there were more than 25,000 cases of polio reported each year. Polio vaccination was begun in 1955. By 1960 the number of reported cases had dropped to about 3,000, and by 1979 there were only about 10. The success of polio vaccination in the U.S. and other countries has sparked a world-wide effort to eliminate polio.

Today: Polio has been eliminated from the United States. But the disease is still common in some parts of the world. It would only take one person infected with polio virus coming from another country to bring the disease back here if we were not protected by vaccine. If the effort to eliminate the disease from the world is successful, some day we won't need polio vaccine. Until then, we need to keep getting our children vaccinated.

3 Who should get polio vaccine and when?

IPV is a shot, given in the leg or arm, depending on age. It may be given at the same time as other vaccines.

Children

Children get 4 doses of IPV, at these ages:

- A dose at 2 months
- A dose at 4 months
- A dose at 6-18 months
- A booster dose at 4-6 years

Some "combination" vaccines (several different vaccines in the same shot) contain IPV.

Children getting these vaccines may get one more (5th) dose of polio vaccine. This is not a problem.

Adults

Most adults 18 and older do not need polio vaccine because they were vaccinated as children. But some adults are at higher risk and should consider polio vaccination:

- (1) people traveling to areas of the world where polio is common,
- (2) laboratory workers who might handle polio virus, and
- (3) health care workers treating patients who could have polio.

Adults in these three groups:

- who have **never been vaccinated against polio** should get 3 doses of IPV:
 - Two doses separated by 1 to 2 months, and
 - A third dose 6 to 12 months after the second.
- who have had **1 or 2 doses** of polio vaccine in the past should get the remaining 1 or 2 doses.



It doesn't matter how long it has been since the earlier dose(s).

- who have had **3 or more doses** of polio vaccine in the past may get a booster dose of IPV.

Your doctor can give you more information.

4

Some people should not get IPV or should wait.

These people should not get IPV:

- Anyone with a life-threatening allergy to any component of IPV, including the antibiotics neomycin, streptomycin or polymyxin B, should not get polio vaccine. Tell your doctor if you have any severe allergies.
- Anyone who had a severe allergic reaction to a previous polio shot should not get another one.

These people should wait:

- Anyone who is moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting polio vaccine. People with minor illnesses, such as a cold, may be vaccinated.

Ask your doctor for more information.

5

What are the risks from IPV?

Some people who get IPV get a sore spot where the shot was given. IPV has not been known to cause serious problems, and most people don't have any problems at all with it.

However, any medicine could cause a serious side effect, such as a severe allergic reaction or even death. The risk of polio vaccine causing serious harm is extremely small.

6

What if there is a moderate or severe problem?

What should I look for?

- Look for any unusual condition, such as a serious allergic reaction, high fever, or unusual behavior.

If a serious allergic reaction occurred, it would happen within a few minutes to a few hours

after the shot. Signs of a serious allergic reaction can include difficulty breathing, weakness, hoarseness or wheezing, a fast heart beat, hives, dizziness, paleness, or swelling of the throat.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS website at www.vaers.hhs.gov or by calling **1-800-822-7967**.

VAERS does not provide medical advice.

7

The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) was created in 1986.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

8

How can I learn more?

- Ask your doctor. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
Polio Vaccine

11/8/2011

42 U.S.C. § 300aa-26

Office Use
Only



Haemophilus Influenzae Type b (Hib) Vaccine

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.


1 What is Hib disease?

Haemophilus influenzae type b (Hib) disease is a serious disease caused by a bacteria. It usually strikes children under 5 years old.

Your child can get Hib disease by being around other children or adults who may have the bacteria and not know it. The germs spread from person to person. If the germs stay in the child's nose and throat, the child probably will not get sick. But sometimes the germs spread into the lungs or the bloodstream, and then Hib can cause serious problems.

Before Hib vaccine, Hib disease was the leading cause of bacterial meningitis among children under 5 years old in the United States. Meningitis is an infection of the brain and spinal cord coverings, which can lead to lasting brain damage and deafness. Hib also cause:

- pneumonia
- severe swelling in the throat, making it hard to breathe
- infections of the blood, joints, bones, and covering of the heart
- death



Before Hib vaccine, about 20,000 children in the United States under 5 years old got severe Hib disease each year and nearly 1,000 people died.

Hib vaccine can prevent Hib disease.

Many more children would get Hib disease if we stopped vaccinating.

2 Who should get Hib vaccine and when?

Children should get Hib vaccine at:

- ✓ 2 months of age
- ✓ 4 months of age
- ✓ 6 months of age*
- ✓ 12-15 months of age

* Depending on what brand of Hib vaccine is used, your child might not need the dose at 6 months of age. Your doctor or nurse will tell you if this dose is needed.

If you miss a dose or get behind schedule, get the next dose as soon as you can. There is no need to start over.

Hib vaccine may be given at the same time as other vaccines.

Older Children and Adults

Children over 5 years old usually do not need Hib vaccine. But some older children or adults with special health conditions should get it. These conditions include sickle cell disease, HIV/AIDS, removal of the spleen, bone marrow transplant, or cancer treatment with drugs. Ask your doctor or nurse for details.

3 Some people should not get Hib vaccine or should wait

- People who have ever had a life-threatening allergic reaction to a previous dose of Hib vaccine should not get another dose.
- Children less than 6 weeks of age should not get Hib vaccine.
- People who are moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting Hib vaccine.

Ask your doctor or nurse for more information.

4

What are the risks from Hib vaccine?


A vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of Hib vaccine causing serious harm or death is extremely small.

Most people who get Hib vaccine do not have any problems with it.

Mild Problems

- Redness, warmth, or swelling where the shot was given (up to 1/4 of children)
- Fever over 101°F (up to 1 out of 20 children)

If these problems happen, they usually start within a day of vaccination. They may last 2-3 days.



5

What if there is a moderate or severe reaction?

What should I look for?

Any unusual condition, such as a serious allergic reaction, high fever or behavior changes. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat, or dizziness within a few minutes to a few hours after the shot.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not provide medical advice

6

The National Vaccine Injury Compensation Program

In the rare event that you or your child has a serious reaction to a vaccine, a federal program has been created to help you pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit the program's website at www.hrsa.gov/vaccinecompensation

7

How can I learn more?

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department's immunization program.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)**
 - Visit the National Immunization Program's website at www.cdc.gov/vaccines



U.S. Department of Health & Human Services
Centers for Disease Control and Prevention



Vaccine Information Statement

Hib (12/16/98)

42 U.S.C. § 300aa-26